

Kids will be kids!

Here are some helpful hints to keep their tummies in check:



Sip on this



When drinking through a straw, many children suck in excess air along with liquid, which can lead to gas. When in doubt, skip the straw!

Take a "time out"



Remind kids to wait a few minutes after eating before running around or playing, to avoid gas and bloating.

Now that's party planning



Parties can be a recipe for overeating—and the indigestion and gas that may result. Have your child choose a few favorite foods and ignore the rest.

When tummy discomfort can't be prevented, turn to **Children's Mylicon**[®] for relief for kids ages 2 through 11

WEIGHT	AGE	Children's Mylicon [®] Tummy Relief Chewables Dosage (Cherry Flavor)	<i>new!</i> Children's Mylicon [®] All-in-One Tummy Relief Liquid Dosage (Bubble Gum Flavor)
Under 24 lbs	Under 2 years	Ask a doctor	Ask a doctor
24-47 lbs	2-5 years	1 tablet – <i>Not more than 3 tablets in 24 hours</i>	5 mL – <i>Not more than 15 mL in 24 hours</i>
48-95 lbs	6-11 years	2 tablets – <i>Not more than 6 tablets in 24 hours</i>	10 mL – <i>Not more than 30 mL in 24 hours</i>

Do not use the maximum dosage for more than 2 weeks except under the advice and supervision of a doctor.

Active ingredients (per tablet or 5 mL liquid dose): calcium carbonate (antacid) 400 mg; simethicone (antigas) 40 mg.

Important: Ask a doctor or pharmacist before use if your child is presently taking a prescription drug. Antacids may interact with certain prescription drugs. See full directions on product package.

To learn more, go to Mylicon.com



Use only as directed.



Scan to buy now!

